

# Javelin

**Javelin** combines the synergies of three active ingredients in combination to give the most effective control on a wide range of broadleaf weeds. Javelin has some residual weed control and can be used early in the season with a pre-germination effect.



**CREEPING OXALIS**



**WHITE CLOVER**



**BINDII**

## OUTSTANDING FEATURES

- No odour.
- Outstanding broad-spectrum weed control.
- Excellent turf safety (cool and warm season grasses).
- S5 Scheduling – safer when handling, mixing and applying.
- Low application rates of active ingredient – safer to the public and the environment.
- Non-ester formulation – reduces potential for off-target damage.
- Contains three active constituents for a wider spectrum of broadleaf weed control.
- Additional surfactants not required.

## IMPORTANT INFORMATION

- Turf clippings should not be used as mulch around other plants or in compost.
- A slight discolouration may occur on kikuyu and carpet grass.
- Due to residual effects of diflufenican overseeding treated areas within three months of application may inhibit establishment of the turf grass.



**globe**  
GROWING SOLUTIONS

# Javelin



**CAPEWEED**



**PLANTAIN**



**CATS EAR**

## **RATE OF APPLICATION**

- Apply at between 4 and 5 L/ha to actively growing weeds.
- Use the lower rate for smaller weeds.
- Complete weed death may take up to six weeks. Faster results can be expected on smaller weeds.

## **MIXING THE PRODUCT**

- Shake the original container before use.
- Agitate continuously to ensure thorough mixing before and during application.
- Mix only sufficient chemical for each days work.
- A surfactant can be added for hard to wet weeds. It should be added after Javelin and water are mixed when the spray tank is almost full.

## **CRITICAL COMMENTS**

- Apply to actively growing weeds.
- Transient discolouration may occur on kikuyu and carpet grass and Queensland blue couch.
- Varietal differences in certain buffalo grasses (eg. ST85) may produce more pronounced effects and it is recommended that small areas be tested for turf safety before largescale application occurs.
- Avoid exceeding application rates through overlapping sprayed areas.
- Use a surfactant for difficult to wet weeds.

## **ACTIVE CONSTITUENTS**

20 g/L clopyralid; 15 g/L diflufenican, 300 g/L MCPA

## **PACKAGING**

Pack size: 10 L

**globe**  
GROWING SOLUTIONS