

# Broadforce MA

## Selective Herbicide

*Broadleaf weed herbicide with excellent safety on lawns*

### Product Overview

BroadForce MA Selective Herbicide is a broadleaf herbicide containing 200g/L MCPA and 200g/L Bromoxynil.

It is registered for the control of a very broad range of dicot weeds in a large range of lawn situations. It is labelled for use on Buffalo, Couch, Qld Blue Couch, Kikuyu, Bent, Fescue, Paspalum and Ryegrass turf types.

BroadForce MA provides quick weed knockdown as well as being one of the broadest spectrum broadleaf weed herbicides available for the Lawn Care market.

It is registered to control the following weeds: Bindii, Jo-Jo, Cudweed, Dandelion, Catsear, Plantains, Thistles, Creeping Oxalis, Capeweed, Creeping Buttercup, Fleabane and Clover.

BroadForce MA is available in 500mL and 1L pack sizes.

### Key Features

- > Safe across a very broad range of lawns, including key varieties such as Buffalo.
- > Registered for the control of 12 common broadleaf weeds.
- > Works on difficult to control weeds such as Creeping Oxalis, Fleabane and Capeweed.
- > Generally controls weeds within 7-21 days following application.
- > Available in two pack sizes – 500mL which treats 800m<sup>2</sup> and 1L which treats 1600m<sup>2</sup>.



Creeping Oxalis

### Maximising performance

**How To Prepare:** Ensure correct dose rate is applied. Half fill the sprayer or watering can with water, add the required amount of product, agitate thoroughly then add the remainder of the water. Do not mix more than is needed. Wash equipment thoroughly after use.

**How To Apply:** Measure lawn area before application. Apply evenly over the area to be treated while weeds are actively growing. Weeds will die slowly; effects may take up to 3 weeks to appear. Repeat applications after 3 to 4 weeks may be necessary to control advanced weeds or certain types of clover. Slight browning of grass may occur under hot conditions.

Important application considerations are as follows:

- > Avoid application during hot or windy conditions, or if rain is expected within 24 hours.
- > Do not apply if shade temperatures are expected to exceed 30°C within a 72 hour period.
- > Avoid application when soil is dry and lawn is suffering from moisture stress.
- > Do not mow or fertilise for 7 days before or after application.

### Broadforce MA Selective Herbicide – Use Rates & Label Recommendations

SITUATION	WEEDS CONTROLLED	MIXING RATE
Established lawns with species as listed: Buffalo, Couch, Qld Blue Couch, Kikuyu, Bent, Fescue, Paspalum, Ryegrass	Bindii, Jo-Jo (Onehunga), Cudweed, Dandelion, Catsear, Plantains, Thistles, Creeping Oxalis, Capeweed, Creeping Buttercup, Fleabane, Clover	Apply evenly over the area to be treated. <b>Sprayer:</b> 30mL per 5-10L of water per 50m <sup>2</sup> <b>Watering Can &amp; Sprinkler Bar:</b> 30mL per 10-20L of water per 50m <sup>2</sup>



Capeweed