



Product Name: CONTRAC BLOX
APVMA Approval No: 48372/125107



Label Name:	CONTRAC BLOX
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	0.05 g/kg BROMADIOLONE Contains Denatonium Benzoate (Bittering Agent)
Mode of Action:	
Statement of Claims:	For the control of rats and mice including those resistant to other anticoagulants, in and around domestic, commercial, public services and industrial buildings.
Net Contents:	NET 1.8 - 9 kg CONTAINS 28 g or 450 g RODENT BLOCKS WHICH ARE ILLEGAL TO SELL SEPARATELY
Restrains:	DO NOT exceed 3 m between bait stations for mice, or 9 m for rats. DO NOT use bait stations further than 2 m from buildings.
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	
Trade Advice:	

General Instructions:	<p>GENERAL INSTRUCTIONS</p> <p>Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoilt bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.</p>
-----------------------	---

Resistance Warning:	
---------------------	--

Precautions:	<p>The use of this product for the control of protected native rodents requires permission from wildlife authorities.</p> <p>DO NOT allow bait to contaminate foodstuffs or food intended for human or animal consumption.</p> <p>DO NOT apply bait to crops.</p> <p>DO NOT place in positions where there is a possibility of contaminating food or food surfaces that come into direct contact with food. The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces.</p>
--------------	--

Protections:	<p>PROTECTION OF WILDLIFE, FISH CRUSTACEANS AND ENVIRONMENT</p> <p>DO NOT contaminate dams, rivers, streams or drains with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.</p>
--------------	---

Storage and Disposal:	<p>Store in the closed, original container in a well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food and feedstuffs, seed and fertilisers. Do not dispose of chemicals on site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product. Dispose of carcasses safely by burning or burying them.</p>
-----------------------	--

Safety Directions:	<p>Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with skin. When preparing product for use wear rubber gloves. If on skin and after each baiting, wash thoroughly with soap and water. Vitamin K1 (Phytomenadione) is antidotal.</p>
--------------------	--

First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26; New Zealand 0800 764 766.</p>
-------------------------	---

First Aid Warnings:	
---------------------	--

DIRECTIONS FOR USE [28 g blocks]

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships) and around grain terminals.	Rats (<i>Rattus norvegicus</i> , <i>Rattus rattus</i>)	2 to 8 x 28 g blocks per bait station	Eliminate as far as practicable, all alternative food sources. Bait areas where conditions might encourage rodent activity. Inspect bait stations on a regular basis. Replace eaten or old/deteriorated baits. If eaten quickly, increase the number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.
	Mice (<i>Mus musculus</i>)	1 to 2 x 28 g blocks per placement or bait station	

DIRECTIONS FOR USE [450 g blocks]

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships) and around grain terminals.	Rats (<i>Rattus norvegicus</i> , <i>Rattus rattus</i>)	1 block per bait station	Eliminate as far as practicable, all alternative food sources. Bait areas where conditions might encourage rodent activity. Inspect bait stations on a regular basis. Replace eaten or old/deteriorated baits. If eaten quickly, increase the number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.
	Mice (<i>Mus musculus</i>)	1 block per infested area	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.