

Product Name: Sundew BattleaxePRO Roach Bait
APVMA Approval No: 82708/110164



Label Name:	Sundew BattleaxePRO Roach Bait
Signal Headings:	READ SAFETY DIRECTIONS
Constituent Statements:	0.5 g/kg FIPRONIL
Mode of Action:	
Statement of Claims:	A crevice, crack or spot treatment for the control of cockroaches.
Net Contents:	5g - 100g
Restraints:	Do not place in areas that are routinely washed.
Directions for Use:	
Other Limitations:	
Withholding Periods:	
Trade Advice:	
General Instructions:	GENERAL INSTRUCTIONS To apply the gel, remove the cap on the nozzle and touch the tip to the surface to be treated and depress plunger slightly. Recap the dispenser after treatment is completed and

return cartridge to outer pack. Re-apply according to remaining level of infestation, when bait is no longer visibly present. A visual inspection of gel placements is recommended one month after initial treatment.

**Resistance
Warning:**

Precautions:

Protections:

PROTECTION OF LIVESTOCK, WILDLIFE AND OTHERS
DO NOT contaminate streams, rivers or waterways with the chemical or used containers.
DO NOT apply this product in any areas easily accessible to children or animals.

**Storage and
Disposal:**

STORAGE AND DIPOSAL
Store in the closed original container in a cool, dry place, out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in a plastic bag and putting in garbage.

Safety Directions:

SAFETY DIRECTIONS
May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

**First Aid
Instructions:**

FIRST AID
If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone 131126 Australia-wide.

First Aid Warnings:

Sundew BattleaxePRO Roach Bait

DIRECTIONS FOR USE

Situation	Pest	Rate	Critical comments
Domestic, commercial and public service buildings	German and other common cockroaches	Apply 1-3 spots per m ² . Recommended spot size is 0.03-0.06 g. Increase the number of spots (up to 3) and spot size depending upon severity of infestation and species present.	Gel placement should be at or near areas where cockroaches gather, such as in cracks and crevices.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION