

LABEL.

David Grays Mancozeb Fungicide 750 DF

DATE: 28 June 2010

Page 1 of 3

(Text appearing above this line does not form part of the label)



CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING**



DAVID GRAYS

Mancozeb Fungicide 750 DF

ACTIVE CONSTITUENT: 750g/kg MANCOZEB

For control of various Fungal Diseases on some Fruit and Vegetables, Flowers and Shrubs in the home garden.

Contents Net: 125 g

APVMA Approval Number: 64791/125g/0410

LABEL.

David Grays Mancozeb Fungicide 750 DF

DATE: 28 June 2010

Page 2 of 3

(Text appearing above this line does not form part of the label)

DAVID GRAYS MANCOZEB FUNGICIDE 750 DF**HOW TO USE:**

- DO NOT apply to if frosts are expected.

Situation	Disease	Rate	How to Apply
BEANS	Rust, Anthracnose	Mix 20g per 10L of water.	Spray at first sign of disease. Repeat at 7-10 day intervals while the risk of disease exists.
CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS	Anthracnose, Downy Mildew		
CARROTS	Leaf Spot		
CARNATIONS, GERANIUMS, HOLLYHOCKS, SNAPDRAGONS, CHRYSANTHEMUMS, GLADIOLI	Rust		
PUMPKINS, MELONS, CUCUMBER, CANTALOUPE SQUASH, ZUCCHINI	Anthracnose, Downy Mildew		
LETTUCE			
MANGOES	Anthracnose		Spray at weekly intervals during flowering then monthly until near harvest.
ONION	Downy Mildew, Purple Blotch		Spray at first sign of disease. Repeat at 7-10 day intervals while the risk of disease exists. DO NOT apply to Tomato seedlings before transplanting.
POME FRUIT: (Apples & Pears)	Black Spot		
ROSES			
TOMATO	Leaf Blight		
TURF	Brown Patch, Helminthosporium Algae	Mix 15g per 10L of water	Apply 10L of mix per 10 square metres of Turf. Repeat every 10 days when disease is present.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED.

LABEL.

David Grays Mancozeb Fungicide 750 DF

DATE: 28 June 2010

Page 3 of 3

(Text appearing above this line does not form part of the label)

WITHHOLDING PERIOD

DO NOT pick carrots, broccoli, cucumber, cantaloupe, pumpkins, melons, zucchini, squash, cabbage, cauliflower, brussel sprouts, onions, beans, for 7 days after spraying.

DO NOT pick apples, pears, tomatoes, lettuce, and mangoes for 14 days after spraying.

HOW TO PREPARE:

Product can be applied through most garden sprayers. Add about half the required volume of water to the sprayer and then add the product. Mix or agitate and then add the remaining water. Regular agitation during spraying is necessary to avoid product settling out. Do not store mixed product. Only mix what you can use immediately. Thoroughly cover plants with spray ensuring all leaves are wetted. Avoid excessive runoff.

CAUTION:

Dangerous to fish. DO NOT allow chemical containers or sprays to get into drains, sewers, streams or ponds. DO NOT feed grass clippings from treated areas to poultry or animals.

STORAGE AND DISPOSAL:

Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS:

May irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. Wash hands after use.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. (Phone 13 11 26). Avoid giving alcohol.

MSDS

Additional information is listed in the Material Safety Data Sheet which is available at www.davidgray.com.au or by contacting David Gray & Co Pty Ltd on (08)9337 4933.

CONDITION OF SALE

David Gray Co. Pty Ltd. shall not be liable for any loss, injury death or damage whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply use or application of this product.

David Gray & Co. Pty. Ltd.
2 Rawlinson Street
O'CONNOR WA 6163
Phone: (08) 9337 4933
DOM:

Batch No:

BARCODE: